

United States Patent [19]

Sogabe et al.

[11] Patent Number:

6,080,553

[45] Date of Patent:

*Jun. 27, 2000

| [54] | CREATINE AMIDINOHYDROLASE, |
|------|----------------------------|
| | PRODUCTION THEREOF AND USE |
| | THEREOF |

[75] Inventors: Atsushi Sogabe; Takashi Hattori;

Yoshiaki Nishiya; Yoshihisa Kawamura, all of Tsuruga, Japan

[73] Assignee: Toyo Boseki Kabushiki Kaisha,

Osaka, Japan

[*] Notice: This patent issued on a continued pros-

ecution application filed under 37 CFR 1.53(d), and is subject to the twenty year patent term provisions of 35 U.S.C.

154(a)(2).

[21] Appl. No.: 08/799,897

[22] Filed: Feb. 13, 1997

[30] Foreign Application Priority Data

[51] Int. Cl.⁷ C12Q 1/34; C12N 9/78; C12N 1/20; C12N 1/00

[56] References Cited

U.S. PATENT DOCUMENTS

3,806,420 4/1974 Holz et al. 195/66

FOREIGN PATENT DOCUMENTS

62 091182 4/1987 Japan . 07 265074 10/1995 Japan .

Primary Examiner—Elizabeth Slobodyansky Attorney, Agent, or Firm—Leydig, Voit & Mayer, Ltd.

] ABSTRACT

A creatine amidinohydrolase having the following physicochemical properties:

Action: catalyzing the following reaction;

creatine+H2O→sarcosine+urea

Optimum temperature: about 40-50° C.

Optimum pH: pH about 8.0-9.0

Heat stability: not more than about 50° C. (pH 7.5, 30

min)

Km value for creatine in a coupling assay using a sarcosine oxidase and a peroxidase: about 3.5-10.0 mM

Molecular weight: about 43,000 (SDS-PAGE)

Isoelectric point: [3.5] 4.5,

a method for producing said enzyme, comprising culture of microorganism producing said enzyme, a method for the determination of creatine or creatinine in a sample using said enzyme, and a reagent therefor.

23 Claims, 2 Drawing Sheets